

# H- $\rightarrow$ bb Note Plans for Summer





Ricardo Gonalo (RHUL)

HSG5 H- $\rightarrow$ bb weekly meeting, 28 June 2011

# News! News! News!



- About  $1.2 \text{ fb}^{-1}$  collected with stable beams so far ( $1.27 \text{ fb}^{-1}$  delivered)
- Peak lumi stable at around  $1.26 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$
- $30 - 50 \text{ pb}^{-1}$  per day (peak so far was  $60 \text{ pb}^{-1}$ )
- 1318 colliding bunches



# News! News! News!

H → WW → lνlν excess


- Excess in data in all channels being investigated
  - See Magda Cheltowska's talk last week: <https://indico.cern.ch/getFile.py/access?contribId=7&resId=0&materialId=slides&confId=143469>
- Cross checks between several analyses, event scans, the works
- Open issues:
  - Problem with the top background
  - Wrong tau polarization in MC
  - Mismodeling of the low p<sub>T</sub> electrons





# News! News! News!

## Statistics Forum

- All ATLAS results for summer conferences should be made using CLs
- (All except analyses with a Bayesian history and Bayesian usage in CMS)



# WH/ZH Note




- Second meeting with the editorial board last week
  - Next meeting right after weekly meeting today
- Note available in CDS:
  - <http://cdsweb.cern.ch/record/1307560?ln=en>
    - List of authors to be updated soon (sorry, mea culpa)
- Our deadlines next...

# Last call for EPS

- **June 23<sup>rd</sup>**: final “analysis update”
    - Assumes ~ 7-10 days to run full analysis
    - **Data taken up to June 18<sup>th</sup> available with GRL (recorded  $L=1 \text{ fb}^{-1}$ );**
  - **Jul 5<sup>th</sup>**: Drafts submitted to EdBoard
    - Early drafts should be circulated to the EdBoard earlier; very similar to PLHC
    - Final drafts on Jul 8<sup>th</sup> to PC;
  - **Jul 9<sup>th</sup>**: last circulation of **Papers/Notes**
    - **Run again the full analysis to include the data after June 26<sup>th</sup> , at least for benchmark analyses? (technical stop on June 29<sup>th</sup>, for ~10 days)**
  - **Jul 15<sup>th</sup>**: last approval meetings
    - Present the circulated results and the ones obtained updating the analysis
  - **Jul 18<sup>th</sup>**: conclude sign-offs of **Papers/CONF Notes**
    - Assumes 1 week for EdBoard and two sign-offs from PC/PubComm/Mngt
    - 18→20: 3 days contingency
  - **Jul 21<sup>st</sup>**: EPS Conference
- 
- This particular schedule can be adopted only for a few Papers and/or Notes (searches or more in general luminosity sensitive)
  - Expect a “hot period” during next two weeks fully dominated by CONF/Paper approval meetings

# Schedule for Setting Limits



- Combination meeting: <https://indico.cern.ch/conferenceDisplay.py?confId=144353>
- Inputs need to be:
  - Available today (28<sup>th</sup> June) – to have **limits** for INT note approval by Ed.board and Higgs WG on 30<sup>th</sup>
  - On 30<sup>th</sup> June – approval and **final decision** on whether to include in SM Higgs combination
  - **Frozen** by 4<sup>th</sup> July for Higgs approval on the 8<sup>th</sup> July
  - **Interpolated** inputs ready right after Higgs WH approval for input to ATLAS SM Higgs combination
  - ATLAS approval on 15<sup>th</sup> July



# JVF-fixed D3PDs

- **Still need analysis cross check!** – cut-flow comparison between different analyses
- SMWZ D3PDs were produced by Haifeng with the bugfix for the Jet Vertex Fraction bug
  - Contain data from run 178044 (22 March 2011) to run 183021 (2 June)
  - Total (data+MC) is 2.7 TB
- Listed in <https://twiki.cern.ch/twiki/bin/view/Main/PrivateD3PDWithJVFFix>
- Transferred to UKI-LT2-RHUL\_LOCALGROUPDISK
  - Should be accessible to everyone; let me know in case of problems
- More samples were produced and now being transferred to grid
- Thanks to Jonas for helping with remaining samples!

ReqID	DataPattern	DestinationSite	Status	NumDatasets All/Subscribed	SummarySize (GB)
34217	user.haifeng.mc10_7TeV.107*.AlpgeJimmyWenuNp*_pt2...	UKI-LT2-RHUL_LOCALGROUPDISK	subscribed	10 / 10	223.3026
34216	user.haifeng.mc10_7TeV.10*.PythiaB_cc*X.merge.NTUP...	UKI-LT2-RHUL_LOCALGROUPDISK	transfer	2 / 2	72.9053
34215	user.haifeng.mc10_7TeV.10*.PythiaB_bb*X.merge.NTUP...	UKI-LT2-RHUL_LOCALGROUPDISK	subscribed	3 / 3	214.9836
34212	user.haifeng.mc10_7TeV.105200.T1_McAtNlo_Jimmy.mer...	UKI-LT2-RHUL_LOCALGROUPDISK	transfer	3 / 3	57.4581
34211	user.haifeng.mc10_7TeV.116590.WH115Inubb_pythia.me...	UKI-LT2-RHUL_LOCALGROUPDISK	subscribed	1 / 1	2.7188
34210	user.haifeng.data11_7TeV.001*.physics_Muons.merge....	UKI-LT2-RHUL_LOCALGROUPDISK	transfer	67 / 67	946.4857
34152	user.haifeng.data11_7TeV.001*.physics_Egamma.merge...	UKI-LT2-RHUL_LOCALGROUPDISK	transfer	61 / 61	1125.8446
34151	user.haifeng.data11_7TeV.001*.physics_Egamma.merge...	UKI-LT2-RHUL_LOCALGROUPDISK	transfer	4 / 4	51.8227
34150	user.haifeng.data11_7TeV.00178109.physics_Egamma.m...	UKI-LT2-RHUL_LOCALGROUPDISK	subscribed	1 / 1	18.7077
34148	user.haifeng.data11_7TeV.00178047.physics_Egamma.m...	UKI-LT2-RHUL_LOCALGROUPDISK	done	1 / 1	0.4138
34147	user.haifeng.data11_7TeV.00178044.physics_Egamma.m...	UKI-LT2-RHUL_LOCALGROUPDISK	done	1 / 1	15.2501



- Submitted abstract to Lepton Photon 2011 in Mumbai, starting August 22<sup>nd</sup>
  - ATLAS  
abstracts:<https://indico.cern.ch/conferenceDisplay.py?confId=143052>
- If accepted, will be presented by Koloina Randrianarivony who volunteered for this

### H->bb searches with the ATLAS detector at the LHC

The H -> bb channel is extremely important for the observation of a Higgs boson signal at the LHC. In the Standard Model, this channel would provide a significant contribution to the Higgs boson search in the low mass region, where this decay mode constitutes the dominant Higgs decay channel. Due to the enormous jet production cross-section at the LHC, the search must target channels where the Higgs boson is produced in association with a weak boson, a pair of top quarks, or jets separated by a rapidity gap. It also requires complex techniques to reconstruct the signal and separate it from an overwhelmingly large background. We present the status of Higgs searches in the H->bb channel currently being performed within ATLAS.

# Other issues...

- PAT group has a vision...
  - See Karsten Koeneke's talk in ATLAS Week


## **What do we want to achieve?**

### **Enable physics analysis in a consistent and efficient manner**

#### **For that to be possible, we need to promote a baseline analysis model**

- As much as possible, every (class of) analysis should start with an Athena-based analysis that results in a dedicated, very specific and small DAOD/D3PD
  - Possibly through intermediate DAODs
  - Lower wall-clock time to iterate on the DAOD/D3PD
  - Possible to store everything locally
- Provide a fast and efficient way to read in the final DAOD/D3PD

# Backup



# WH/ZH Note: Missing Ingredients

- Moving to MC10b: done
- b tagging:
  - Need advanced tagger for increased background rejection
  - Efficiency scale factors almost done
  - Calibration & fake rate: preliminary on week of 20th June - will re-do analysis with final numbers
  - IP3D+SV1, 60% efficiency working point
- Jet Vertex Fraction:
  - Fix exists but applicable only to AOD-based analyses – i.e. only one analysis in our group
  - D3PDs including the bug fix exist for part of the 2011 run – being transferred to Grid site
  - **Validation?**
- Editorial board:
  - Richard Bateley (chair)
  - Alex Read
  - Emmanuel Lemonier
  - Niels van Eldik
  - Good 1<sup>st</sup> meeting with Ed.Board
- QCD background (incl. bb, cc):
  - Almost there
  - One **feature to be understood** in anti-track isolation QCD background in electron channel
  - Watch this space!
- Systematics:
  - First estimates done – dominated by b-tagging uncertainty (around 30%)
- SM Higgs combination:
  - Need to produce inputs for SM Higgs combination

# LHC actual versus design parameters


	design	present	comment
Beam energy	7 TeV	<b>3.5 TeV</b>	½ design
transv. norm. emittance	3.75 $\mu\text{m}$	<b>2.9 <math>\mu\text{m}</math></b>	¾ design!
beta*	0.55 m	1.5 m	3x design
IP beam size	16.7 $\mu\text{m}$	34 $\mu\text{m}$	2x design
bunch intensity	1.15x10 <sup>11</sup>	<b>1.25x10<sup>11</sup></b>	higher than design
luminosity / bunch	3.6x10 <sup>30</sup> cm <sup>-2</sup> s <sup>-1</sup>	1.1x10 <sup>30</sup> cm <sup>-2</sup> s <sup>-1</sup>	only factor 3 away (x4 from energy!)
# bunches	2808	1092	approaching ½ design
bunch spacing	25 ns	<b>50 ns</b>	
beam current	0.582 A	0.236 A	close to ½ design
rms bunch length	7.55 cm	≥8.7 cm	
crossing angle	285 $\mu\text{rad}$	240 $\mu\text{rad}$	
“Piwinski angle”	0.64	≥0.31	
luminosity	10 <sup>34</sup> cm <sup>-2</sup> s <sup>-1</sup>	1.2x10 <sup>33</sup> cm <sup>-2</sup> s <sup>-1</sup>	>10% design

# Do we need a JVF cut?

- In principle yes!...
- Need to use cut  $N_{\text{jets}} = 2$  to suppress  $t\bar{t}$  background; use  $N_{\text{jets}} = 3$  as  $t\bar{t}$  control region
- So must suppress spurious jets from pileup...


$N_{\text{jets}} < 4$

All backgrounds from Monte Carlo  
bb and cc MC clearly not enough



$N_{\text{jets}} = 2$

QCD background from data  
Before last scale factor (1-b sideband)






# Do we need a JVF cut?

- In fact, not using the Jet Vertex Fraction seems to have a significant effect on Njets
- But a small effect after all cuts...

	data	tt MC
No JVF cut	303	200
JVF > 0.75	300	185


Haifeng Li

With JVF



Haifeng Li

No JVF





# WH/ZH Note: Outlook

- Skeleton draft of INT note should be available now...
- Then a couple of weeks to finish details of QCD BG determination and interact with Editorial Board
  - Expect some changes to cuts etc during this
- Dataset frozen on 22 June (I think)
- Preliminary b-tagging calibrations around same time
- Aim for Higgs approval at end of June
- Last iteration with final b-tagging calibrations on...
- Circulate note to ATLAS for CONF approval in early July for approval in time for EPS